

Dynamic Causal Regression for Functional Data

Daniel Gervini

Department of Mathematical Sciences

University of Wisconsin–Milwaukee

January 14, 2013

Abstract

A characteristic feature of samples of curves is the presence of time variability in addition to amplitude variability. Existing functional regression methods do not handle time variability in an efficient way. In this paper we propose a functional regression method that incorporates time warping as an intrinsic part of the model, and then attains high predictive power in a parsimonious way, avoiding overfitting. The properties of the estimators are studied by simulation, and an application to the study of human lip movement is presented.

Key Words: Causal Linear Regression; Functional Data Analysis; Registration; Spline Smoothing; Time Warping.

1 Introduction

Many statistical applications today involve modeling curves as functions of other curves. For example, the evolution of CD4 cell counts in HIV patients can be modeled as a function of viral load (Liang et al. 2003, Wu and Liang 2004, Wu and Müller 2010); gene expression profiles of insects at the pupa stage can be modeled as functions of gene expression profiles at the embryonic stage (Müller et al. 2008); systolic blood pressure trajectories over the years can be predicted to some extent from body mass index trajectories (Yao et al. 2005). These are just a few examples that show the broad applicability of functional regression methods.

Functional regression, as applied in the papers cited above, is a more or less straightforward extension of multivariate linear regression to the functional-data framework (Ramsay and Silverman 2005, ch. 16). Recent developments have focused on theoretical aspects such as rates of convergence (Cai and Hall 2006, Hall and Horowitz 2007, Crambes et al. 2009), sparse longitudinal data (Yao et al. 2005), and interpretability of the estimators (James et al. 2009). But a problem inherent to functional data that has received little attention in the regression context is the problem of time variability.

As a motivating example, consider the problem in Malfait and Ramsay (2003). The authors wanted to predict lip acceleration using electromyography (EMG) curves that measure neural activity in the primary muscle depressing the lower lip, the depressor labii inferior. A person was asked to repeat the phrase “say Bob again” a few times, and the lip movement and associated EMG curve corresponding to the word “Bob” were recorded. Lip acceleration curves were obtained by differentiating the smoothed lip trajectories. The sample curves, time-standardized to 700 msec, are shown in Figure 1. Both types of curves follow regular patterns, but they show considerable variability both in amplitude and timing of the main features. In fact, time variability overwhelms amplitude variability in Figure 1(a), to the point that it is not clear whether a typical EMG curve has three peaks or four. A pair-by-pair analysis of the curves shows that the EMG spikes are well aligned with certain features of the acceleration curves; therefore, the *timing* of the EMG spikes (not just their amplitude) is likely to provide valuable information for predicting lip acceleration.

The classical functional linear regression model does not explicitly model the dynamics of the curves; instead, it sees time variability as a nuisance. Time variability tends to spread the features of both the predictor and the response curves over wide time ranges, with the result that the regression function needs to be very irregular to provide a good fit to the data; overfitting may become a problem for small sample sizes, like those in Figure

Figure 1: Lip Movement Example. (a) EMG curves; (b) lip acceleration curves.

1 (which consists of only 29 curves). On the other hand, if the curves are well aligned, a smoother and more localized regression function will be sufficient to provide a good fit to the data. The situation is similar to the problem of estimating the Karhunen–Loève components: although in theory any squared-integrable sample can be approximated by a linear combination of orthogonal functions, in practice it is more efficient to remove as much of the time variability as possible first by synchronizing the data, and then to estimate the (fewer) Karhunen–Loève components of the aligned curves. For a more general discussion of the registration problem see Ramsay and Silverman (2005, ch. 7) and Kneip and Ramsay (2008).

A similar approach could be attempted for the regression problem; that is, to align the curves first and then fit a regression model to the synchronized curves. There are many warping methods available for this, such as Gervini and Gasser (2004, 2005), James (2007), Kneip et al. (2000), Liu and Müller (2004), Ramsay and Li (1998), Tang and Müller (2008, 2009), and Wang and Gasser (1999). But the registration of the covariates and the responses would need to be carried out independently, and it would not be possible to predict the latter from the former, unless a parallel regression model is also set up for the warping functions. This would be cumbersome and statistical inference would be complicated.

Given that time variability cannot be ignored but there are no methods that handle time variability efficiently in a regression context, we propose in this paper a regression method that incorporates time warping as an intrinsic part of the model. Since we are going to apply this method to the above lip movement data, we will focus on the causal regression model, or “historical linear model” as Malfait and Ramsay (2003) call it; that is, the model does not use the whole trajectory of a covariate to predict a response value

at time t , but only those observations preceding time t . In addition, we will assume that the sample curves are smooth curves; in effect, we are assuming that the raw data has been pre-processed by smoothing. Incorporating warping into the traditional (non-causal) functional regression model and modeling sparse and irregular trajectories are important problems but they will be addressed in other papers.

This paper is organized as follows. The dynamic causal regression model is introduced in Section 2. A convenient semiparametric family of warping functions is discussed in Section 3. Asymptotics and inference are discussed in Section 4. The finite-sample behavior of the estimator is studied by simulation in Section 5. Finally, the lip movement data mentioned above is analyzed in more detail in Section 6. Matlab[®] programs implementing these methods are available on the author's website.

2 Dynamic causal regression

2.1 The model

Let $(x_1, y_1), \dots, (x_n, y_n)$ be a sample of functions, where x_i is the covariate and y_i the response. We assume x_i and y_i are square-integrable functions on a the same time interval $[a, b]$. A linear predictor of $y_i(t)$ based on x_i has the form

$$L(t; x_i, \alpha, \beta) = \alpha(t) + \int_a^b \beta(s, t) x_i(s) ds, \quad (1)$$

where α is the intercept function and β the slope function. However, (1) employs the whole trajectory $\{x_i(s) : s \in [a, b]\}$ to predict the value of y_i at t , including “future” observations $x_i(s)$ with $s > t$. In many applications this is not reasonable. For example, for the lip movement data in Figure 1 it is clear that future neural activity cannot have an influence on past lip movement; therefore, prediction of $y_i(t)$ must be based only on the partially observed curves $\{x_i(s) : s \in [a, t]\}$. We incorporate this restriction into model (1) by assuming $\beta(s, t) = 0$ for $s > t$. This model is called “historical linear model” by Malfait and Ramsay (2003), although we prefer to call it “causal linear regression”.

As explained in Section 1, linear regression works best for synchronized curves. Suppose, then, that for each pair (x_i, y_i) we have a warping function $w_i : [a, b] \rightarrow [a, b]$, strictly monotone increasing and such that $w_i(a) = a$ and $w_i(b) = b$. Let $\tilde{x}_i = x_i \circ w_i$ and $\tilde{y}_i = y_i \circ w_i$ be the warped curves; then we can apply (1) to $(\tilde{x}_i, \tilde{y}_i)$ rather than (x_i, y_i) , and

the *dynamic functional predictor* of $y_i(t)$ is defined as

$$L(w_i^{-1}(t); \tilde{x}_i, \alpha, \beta) = \alpha(w_i^{-1}(t)) + \int_a^b \beta(s, w_i^{-1}(t)) x_i(w_i(s)) ds. \quad (2)$$

Note that the same warping function w_i is used for x_i and y_i ; this is reasonable for the type of applications we have in mind. In particular, using a common warping function preserves the causality of the model: if $\beta(s, t) = 0$ for $s > t$, the integral in (2) only uses those values of $x_i(w_i(s))$ with $s \leq w_i^{-1}(t)$, which is to say only the values of $x_i(u)$ with $u \leq t$.

The estimators of α , β , and the w_i s can be obtained by least squares: if $\|f\| := \{\int_a^b f^2(t) dt\}^{1/2}$ is the usual L^2 norm, we minimize

$$\sum_{i=1}^n \|y_i \circ w_i - L(\cdot; x_i \circ w_i, \alpha, \beta)\|^2 \quad (3)$$

subject to the identifiability condition $\sum_{i=1}^n w_i(t)/n \equiv t$. For given β and w_i s, the α that minimizes (3) is clearly

$$\hat{\alpha}(t) = \bar{y}(t) - \int \hat{\beta}(s, t) \bar{x}(s) ds, \quad (4)$$

so we only need to focus on the estimation of β and the w_i s. For the warping functions we will use the semiparametric family described in Section 3. For β we will also use a semiparametric family: truncated tensor splines. Specifically, if $\{\psi_j(s)\}$ is a (univariate) B-spline basis, we define $\xi_{ij}(s, t) = \psi_i(s)\psi_j(t)I\{s \leq t\}$, drop the ξ_{ij} s that are zero everywhere, and re-scale the remaining p functions so that $\sum_{k=1}^p \phi_k(s, t) \equiv 1$. Then any β of the form $\beta(s, t) = \sum_{k=1}^p b_k \phi_k(s, t)$ satisfies the causality restriction $\beta(s, t) = 0$ for $s > t$. Other bases are possible, such as the simplicial basis used in Malfait and Ramsay (2003).

Typically the dimension p of this space will be large and then the regularity of β must be controlled by some form of roughness penalty. In the smoothing literature it is common to penalize the curvature of the functions, that is, the L^2 norm of the second derivative, which for β would translate into $\iint \|H\beta(s, t)\|_F^2 ds dt$, where H is the Hessian and $\|\cdot\|_F$ the Frobenius norm. However, in the regression context it is more natural to penalize the size of β , as it is done for instance in ridge regression, so we will use $\iint \beta^2(s, t) ds dt$ as

penalty and minimize

$$\sum_{i=1}^n \|y_i \circ w_i - L(\cdot; x_i \circ w_i, \alpha, \beta)\|^2 + \lambda \iint \beta^2(s, t) ds dt. \quad (5)$$

The penalization parameter λ can be chosen via Akaike-like criteria, discussed below.

2.2 Estimation

At this point it is convenient to introduce some notation. If f and g are two square-integrable functions on $[a, b]$, the usual inner product is defined as $\langle f, g \rangle := \int_a^b f(t)g(t) dt$ (for bivariate functions such as $\beta(s, t)$ it is defined analogously.) If now $\mathbf{f} : [a, b] \rightarrow \mathbb{R}^p$ and $\mathbf{g} : [a, b] \rightarrow \mathbb{R}^q$ are vector-valued functions with square-integrable elements, we denote by $\langle \mathbf{f}, \mathbf{g}^T \rangle$ the $p \times q$ matrix with elements $\langle f_i, g_j \rangle$ (in other words, the integration is carried out in a componentwise manner.) With this notation, $(\langle \mathbf{f}, \mathbf{g}^T \rangle)^T = \langle \mathbf{g}, \mathbf{f}^T \rangle$ and $\mathbf{A} \langle \mathbf{f}, \mathbf{g}^T \rangle = \langle \mathbf{A} \mathbf{f}, \mathbf{g}^T \rangle$ for any matrix $\mathbf{A}$ of dimensions compatible with $\mathbf{f}$. Then the penalized least-squares problem (5) can be rewritten as

$$\min_{\mathbf{b}, \{w_i\}} \frac{1}{n} \sum_{i=1}^n \|\tilde{y}_i^* - \mathbf{b}^T \mathbf{z}_i\|^2 + \lambda \mathbf{b}^T \mathbf{\Omega} \mathbf{b}, \quad (6)$$

where $\tilde{y}_i^* := \tilde{y}_i - \bar{\tilde{y}}$, $\tilde{x}_i^* := \tilde{x}_i - \bar{\tilde{x}}$, $\mathbf{z}_i(t) = \langle \phi(\cdot, t), \tilde{x}_i^* \rangle$, and $\mathbf{\Omega} = \langle \phi, \phi^T \rangle$, with $\phi(s, t) = (\phi_1(s, t), \dots, \phi_p(s, t))^T$.

The minimization problem (6) is not quadratic in $\mathbf{b}$ due to the dependence of $\tilde{x}_i^*$ and $\tilde{y}_i^*$ on the warping functions, but it is conditionally quadratic for given warping functions. Therefore we propose an alternating least-squares algorithm: (i) given warping functions $\{w_i\}$, find the $\mathbf{b}$ that minimizes (6), which is simply the ridge estimator

$$\hat{\mathbf{b}}_\lambda = \left(\sum_{i=1}^n \langle \mathbf{z}_i, \mathbf{z}_i^T \rangle + n\lambda \mathbf{\Omega} \right)^{-1} \sum_{i=1}^n \langle \mathbf{z}_i, \tilde{y}_i^* \rangle; \quad (7)$$

(ii) given $\mathbf{b}$, for each $i = 1, \dots, n$ find the $\hat{w}_i$ that minimizes $\|\tilde{y}_i^* - \mathbf{b}^T \mathbf{z}_i\|^2$ over the family of warping functions discussed in Section 3. Steps (i) and (ii) are alternated until convergence. Note that step (ii) by itself would require many iterations if the exact minimizer $\hat{w}_i$ were to be computed, but to save time we recommend to perform just one step towards the solution on each iteration.

Once the estimators $\hat{\alpha}$, $\hat{\beta}$ and $\hat{w}_1, \dots, \hat{w}_n$ have been obtained, it is possible to use (2) to predict a response function y_{n+1} for a new covariate function x_{n+1} . The natural predictor

of y_{n+1} given x_{n+1} is $\hat{y}_{n+1}(t) = L(\hat{w}_{n+1}^{-1}(t); \tilde{x}_{n+1}, \hat{\alpha}, \hat{\beta})$, but $\hat{w}_{n+1}$ cannot be obtained by minimizing (6) because that would involve the unobserved response y_{n+1} . Instead, $\hat{w}_{n+1}$ can be obtained from x_{n+1} alone by registering x_{n+1} to the mean of the warped x_i s:

$$\hat{w}_{n+1} = \arg \min_w \|x_{n+1} \circ w - \hat{\mu}_{\tilde{x}}\|^2, \quad (8)$$

where $\hat{\mu}_{\tilde{x}} = n^{-1} \sum_{i=1}^n \tilde{x}_i$. Note that $\hat{\mu}_{\tilde{x}}$ is fixed here, so the “pinching” or “overwarping” problem associated with least-squares registration [discussed in Ramsay and Silverman (2005, ch. 7.6) and Kneip and Ramsay (2008)] will not be a serious issue.

2.3 Criteria for penalty parameter selection

A crucial issue for the good performance of the estimator is selection of the penalty parameter λ . In the literature this is typically done by cross-validation or by an Akaike-type criterion [see e.g. Hastie et al. (2009)]. A variant of Akaike’s criterion that we found particularly useful is the “corrected Akaike criterion” or AICC, proposed by Hurvitz et al. (1998). To the best of the author’s knowledge, functional-data equivalents of the AICC criterion and of the notion of “effective degrees of freedom” (Hastie et al., 2009) have not yet been derived in the literature, so we do it in detail in this section.

To begin with, we need a functional equivalent of the “hat matrix”, an operator $\mathcal{H} : \{L^2([a, b])\}^n \rightarrow \{L^2([a, b])\}^n$ such that $\widehat{\mathbf{y}}^*(t) = (\mathcal{H}\mathbf{y}^*)(t)$. Let $\mathbf{Z}(t)$ denote the $n \times p$ matrix with rows $\mathbf{z}_i^T(t)$. Then from (7) we have

$$\widehat{\mathbf{y}}^*(t) = \mathbf{Z}(t)\hat{\mathbf{b}}_\lambda = \mathbf{Z}(t)\mathbf{A}_\lambda^{-1} \sum_{i=1}^n \langle \mathbf{z}_i, \tilde{y}_i^* \rangle,$$

where $\mathbf{A}_\lambda = \sum_{i=1}^n \langle \mathbf{z}_i, \mathbf{z}_i^T \rangle + \lambda \mathbf{\Omega}$. Note that, with our notation, $\sum_{i=1}^n \langle \mathbf{z}_i, \tilde{y}_i^* \rangle = \langle \mathbf{Z}^T, \tilde{\mathbf{y}}^* \rangle$ and $\sum_{i=1}^n \langle \mathbf{z}_i, \mathbf{z}_i^T \rangle = \langle \mathbf{Z}^T, \mathbf{Z} \rangle =: \mathbf{A}$. Then the “hat operator” $\mathcal{H}$ is given by $(\mathcal{H}\mathbf{f})(t) = \mathbf{Z}(t)\mathbf{A}_\lambda^{-1} \langle \mathbf{Z}^T, \mathbf{f} \rangle$, where $\mathbf{f}$ is an n -dimensional vector of square-integrable functions.

The hat operator $\mathcal{H}$ is an integral operator: we can write $(\mathcal{H}\mathbf{f})(t) = \int_a^b \mathbf{R}(s, t) \mathbf{f}(s) ds$, with $\mathbf{R}(s, t) = \mathbf{Z}(t)\mathbf{A}_\lambda^{-1} \mathbf{Z}^T(s)$. The trace of an integral operator is defined as $\int_a^b \text{tr}\{\mathbf{R}(s, s)\} ds$, which like its finite-dimensional analogue is equal to the sum of the eigenvalues (Gohberg et al., 2003). For our particular operator $\mathcal{H}$ it is easy to verify that $\text{tr}(\mathcal{H}) = \text{tr}(\mathbf{A}_\lambda^{-1} \mathbf{A})$, so no computations beyond those required for $\hat{\mathbf{b}}_\lambda$ are involved.

Following Hastie et al. (2009) we define $\text{df}(\lambda)$, the “effective degrees of freedom” of the model, as $\text{tr}(\mathcal{H})$. And following Hurvitz et al. (1998) we define the “corrected Akaike

criterion” as

$$\text{AICC}(\lambda) = \text{MSE}(\lambda) \exp[1 + 2\{\text{df}(\lambda) + 1\}/\{n - \text{df}(\lambda) - 2\}],$$

where $\text{MSE}(\lambda) = n^{-1} \sum_{i=1}^n \|\tilde{y}_i^* - \hat{\tilde{y}}_i^*\|^2$. We can also define a functional equivalent of the “generalized cross validation” criterion of Wahba (1990):

$$\text{GCV}(\lambda) = \text{MSE}(\lambda)/\{1 - \text{df}(\lambda)/n\}^2,$$

but this criterion generally leads to undersmoothing, so we will use $\text{AICC}(\lambda)$ in this paper. The performance of $\text{AICC}(\lambda)$ is studied by simulation in Section 5.

3 A semiparametric family of warping functions

The warping functions w_i can also be modeled semiparametrically, but due to the monotonicity restriction some care must be taken in the choice of basis functions. We found that a convenient family are the cubic Hermite splines [see e.g. Fritsch and Carlson (1980)]. Details are given in the technical supplement, but the basic idea is the following. Let $\boldsymbol{\tau} = (\tau_0, \tau_1, \dots, \tau_{r+2})$ be a vector of knots in $[a, b]$, with $a = \tau_0 < \tau_1 < \dots < \tau_{r+2} = b$. The knots can be chosen arbitrarily (for example, equispaced point in $[a, b]$), or they can be made to correspond, roughly, to the average location of the salient features of the curves. For example, in Figure 1(a) one would choose $\boldsymbol{\tau} = (0, .08, .2, .5, .69)$ or perhaps $\boldsymbol{\tau} = (0, .08, .2, .4, .5, .69)$. A cubic Hermite spline can be expressed as $w_i(t) = \sum_{k=0}^{r+1} c_{ik} \eta_k(t; \boldsymbol{\tau}) + \sum_{k=0}^{r+1} d_{ik} \xi_k(t; \boldsymbol{\tau})$, with

$$\eta_k(t; \boldsymbol{\tau}) = \begin{cases} h_0\left(\frac{t-\tau_0}{\tau_1-\tau_0}\right) & \text{if } k = 0 \\ h_0\left(\frac{t-\tau_k}{\tau_{k+1}-\tau_k}\right) + h_0\left(\frac{\tau_k-t}{\tau_k-\tau_{k-1}}\right) - I_{\{\tau_k\}}(t) & \text{if } k = 1, \dots, r \\ h_0\left(\frac{\tau_{r+1}-t}{\tau_{r+1}-\tau_r}\right) & \text{if } k = r + 1, \end{cases}$$

for $h_0(t) = (1 + 2t)(1 - t)^2 I_{[0,1]}(t)$, and

$$\xi_k(t; \boldsymbol{\tau}) = \begin{cases} h_1\left(\frac{t-\tau_0}{\tau_1-\tau_0}\right) (\tau_1 - \tau_0) & \text{if } k = 0 \\ h_1\left(\frac{t-x_k}{\tau_{k+1}-\tau_k}\right) (\tau_{k+1} - \tau_k) - h_1\left(\frac{x_k-t}{\tau_k-\tau_{k-1}}\right) (\tau_k - \tau_{k-1}) & \text{if } k = 1, \dots, r \\ -h_1\left(\frac{\tau_{r+1}-t}{\tau_{r+1}-\tau_r}\right) (\tau_{r+1} - \tau_r) & \text{if } k = r + 1, \end{cases}$$

for $h_1(t) = t(1 - t)^2 I_{[0,1]}(t)$.

Note that $\eta_k(\tau_k; \boldsymbol{\tau}) = 1$ and $\eta_k(\tau_j; \boldsymbol{\tau}) = 0$ for $j \neq k$, while $\xi_k(\tau_j; \boldsymbol{\tau}) = 0$ for all j , so $w_i(\tau_k) = c_{ik}$. Then the coefficients c_{ik} are interpretable: if the τ_k s are average “landmarks”, the c_{ik} s would be the corresponding landmarks of the i th curve, and the warping function w_i is implicitly performing a “landmark registration”. It is also easy to verify that $w'_i(\tau_k) = d_{ik}$. It can be shown that for *any* coefficient vector $\mathbf{c}_i$ with $a = c_{i0} < c_{i1} < \dots < c_{i,r+1} = b$, a set of derivatives d_{ik} can be constructed so that $w_i(t)$ is strictly increasing in $[a, b]$. Fritsch and Carlson (1980) give necessary and sufficient conditions for this, and present a simple one-pass algorithm to construct a set of d_{ik} s that satisfy the sufficient conditions. We will use Fritsch-Carlson algorithm in this paper, so the d_{ik} s will not be free parameters but functions of the c_{ik} s. Then $w_i(t)$ is entirely parameterized by $\mathbf{c}_i$ for a given knot sequence $\boldsymbol{\tau}$.

For computational purposes it is convenient to transform the constrained monotone vector $(c_{i1}, \dots, c_{ir})$ into an unconstrained vector $\boldsymbol{\theta}_i$ via the (invertible) Jupp transform (Jupp, 1978): $\theta_{ik} = \log\{(c_{i,k+1} - c_{ik})/(c_{ik} - c_{i,k-1})\}$, for $k = 1, \dots, r$. The minimization problem in step (ii) of the algorithm presented in Section 2 then becomes a simple unconstrained r -dimensional minimization problem, albeit a non-linear one.

Finally, we note that although we treat $\boldsymbol{\tau}$ as a fixed sequence of knots in this paper, it would not be unreasonable to treat $\boldsymbol{\tau}$ as a free parameter and estimate it as well. The dependence of the basis functions $\{\eta_k\}$ and $\{\xi_k\}$ on $\boldsymbol{\tau}$ is not complicated, and explicit formulas for their derivatives with respect to the knots are easy to obtain, so a Newton-Raphson algorithm to find the optimal $\boldsymbol{\tau}$ could be implemented. However, to keep the paper focused, we will not pursue the free-knot alternative here.

4 Asymptotics

There are two approaches to the asymptotic analysis of dynamic regression estimators. The “nonparametric” approach would investigate the rate of convergence of $\hat{\beta}_n(s, t)$ to the true regression function $\beta_0(s, t)$ when the number of basis functions p increases as n goes to infinity, using techniques as in e.g. Van de Geer (2000, ch. 10); but this is not what we are interested in in this paper. We will follow a “parametric” approach: we will hold p fixed and make the smoothing parameter λ_n go to zero as n goes to infinity [as in Knight and Fu (2000)]. Then the basis coefficients $\hat{\mathbf{b}}_n$ will converge at the usual $n^{1/2}$ rate to a $\mathbf{b}^*$ such that $\beta^*(s, t) = \mathbf{b}^{*T} \boldsymbol{\phi}(s, t)$, which in general will not be the true $\beta_0(s, t)$ but hopefully will be close to it. This simplification ignores estimation bias but allows us to derive a simple estimator of the asymptotic variance of $\hat{\beta}_n(s, t)$.

Let us first introduce some notation. The warping functions are of the form $w(t, \boldsymbol{\theta}_i)$,

with $\boldsymbol{\theta}_i \in \mathbb{R}^r$ an unconstrained parameter. The minimization (6) is carried out simultaneously on $\mathbf{b}$ and the $\boldsymbol{\theta}_i$ s, but for the asymptotic analysis it is best to see (6) as a profile least squares problem: let

$$r_i(\boldsymbol{\theta}, \mathbf{b})(t) = y_i \circ w(t, \boldsymbol{\theta}) - \mu_y(t) - \int_a^b \mathbf{b}^T \boldsymbol{\phi}(s, t) \{x_i \circ w(s, \boldsymbol{\theta}) - \mu_x(s)\} ds,$$

then for each $\mathbf{b}$ define

$$\boldsymbol{\theta}_i(\mathbf{b}) := \arg \min_{\boldsymbol{\theta}} \|r_i(\boldsymbol{\theta}, \mathbf{b})\|^2 \quad (9)$$

and

$$f_i(\mathbf{b})(t) = r_i(\boldsymbol{\theta}_i(\mathbf{b}), \mathbf{b})(t). \quad (10)$$

Then

$$\hat{\mathbf{b}}_n = \arg \min_{\mathbf{b}} \frac{1}{n} \sum_{i=1}^n \|f_i(\mathbf{b})\|^2 + \lambda_n \mathbf{b}^T \boldsymbol{\Omega} \mathbf{b}. \quad (11)$$

For simplicity, we are going to assume μ_x and μ_y are fixed and known. As $n \rightarrow \infty$ and $\lambda_n \rightarrow 0$, $\hat{\mathbf{b}}_n$ will converge in probability to

$$\mathbf{b}^* = \arg \min_{\mathbf{b}} E_0 \{\|f_i(\mathbf{b})\|^2\},$$

where E_0 denotes expectation with respect to the true model:

$$\begin{aligned} \tilde{y}(t) &= \mu_{\tilde{y}}(t) + \int_a^b \beta_0(s, t) \{\tilde{x}(s) - \mu_{\tilde{x}}(s)\} ds + \varepsilon(t), \\ (x(t), y(t)) &= (\tilde{x} \circ w^{-1}(t), \tilde{y} \circ w^{-1}(t)), \end{aligned} \quad (12)$$

for a warping process $w(t)$ and a regression slope $\beta_0(s, t)$ that may not necessarily belong to the families used for estimation. If w and β_0 do belong to the families used for estimation, say $\beta_0(s, t) = \mathbf{b}_0^T \boldsymbol{\phi}(s, t)$, one expects $\mathbf{b}^*$ to be close to $\mathbf{b}_0$, but $\mathbf{b}^*$ will be equal to $\mathbf{b}_0$ only in the ideal case that $\varepsilon(t) \equiv 0$. This is an unfortunate drawback of least-squares registration methods: they need exact specification of the model in order to be consistent [see e.g. Proposition 1 of Kneip and Ramsay (2008)].

Theorem 1 *If $\sqrt{n}\lambda_n \rightarrow \ell_0$ as $n \rightarrow \infty$, then $\sqrt{n}(\hat{\mathbf{b}}_n - \mathbf{b}^*) \xrightarrow{D} N(-\ell_0 \boldsymbol{\Gamma}^{-1} \boldsymbol{\Omega} \mathbf{b}^*, \boldsymbol{\Gamma}^{-1} \boldsymbol{\Lambda} \boldsymbol{\Gamma}^{-1})$ with*

$$\begin{aligned} \boldsymbol{\Gamma} &= E_0 \{\langle \dot{\mathbf{f}}_i(\mathbf{b}^*), \dot{\mathbf{f}}_i(\mathbf{b}^*)^T \rangle\}, \\ \boldsymbol{\Lambda} &= E_0 \{\langle f_i(\mathbf{b}^*), \dot{\mathbf{f}}_i(\mathbf{b}^*) \rangle \langle f_i(\mathbf{b}^*), \dot{\mathbf{f}}_i(\mathbf{b}^*)^T \rangle\}, \end{aligned}$$

where $\dot{\mathbf{f}}_i(\mathbf{b})(t)$ is the gradient of $f_i(\mathbf{b})(t)$ with respect to $\mathbf{b}$.

The proof of Theorem 1 is given in the Appendix. A more explicit expression for $\dot{\mathbf{f}}_i(\mathbf{b})$ can be obtained as follows. Let D_θ and D_b denote the differentials with respect to the variables θ and $\mathbf{b}$, so $\dot{\mathbf{f}}_i(\mathbf{b}) = D_b f_i(\mathbf{b})^T$. Then we have

$$D_b f_i(\mathbf{b})(t) = \{D_\theta r_i(\theta_i(\mathbf{b}), \mathbf{b})(t)\} \{D_b \theta_i(\mathbf{b})\} + D_b r_i(\theta_i(\mathbf{b}), \mathbf{b})(t),$$

where

$$D_\theta r_i(\theta, \mathbf{b})(t) = y'_i(w(t, \theta)) D_\theta w(t, \theta) - \int_a^b \mathbf{b}^T \phi(s, t) x'_i(w(s, \theta)) D_\theta w(s, \theta) ds$$

and

$$D_b r_i(\theta, \mathbf{b})(t) = - \int_a^b \phi(s, t)^T \{x_i \circ w(s, \theta) - \mu_x(s)\} ds.$$

From Section 3 we have

$$D_\theta w(t, \theta) = \eta(t; \tau)^T D_\theta \mathbf{J}^{-1}(\theta)$$

with $\eta(t; \tau)^T = (\eta_1(t; \tau), \dots, \eta_r(t; \tau))$ and $\mathbf{J}^{-1}(\theta)$ the inverse Jupp transform. To obtain $D_b \theta_i(\mathbf{b})$ note that (9) implies

$$\langle r_i(\theta_i(\mathbf{b}), \mathbf{b}), D_\theta r_i(\theta_i(\mathbf{b}), \mathbf{b}) \rangle = \mathbf{0}^T \text{ for all } \mathbf{b},$$

so $D_b \theta_i(\mathbf{b})$ can be derived via the Implicit Function Theorem: if $\mathbf{F}_i(\theta, \mathbf{b}) = \langle r_i(\theta, \mathbf{b}), D_\theta r_i(\theta, \mathbf{b})^T \rangle$, then

$$D_b \theta_i(\mathbf{b}) = -\{D_\theta \mathbf{F}_i(\theta_i(\mathbf{b}), \mathbf{b})\}^{-1} \{D_b \mathbf{F}_i(\theta_i(\mathbf{b}), \mathbf{b})\},$$

where

$$D_\theta \mathbf{F}_i(\theta, \mathbf{b}) = \langle D_\theta r_i(\theta, \mathbf{b})^T, D_\theta r_i(\theta, \mathbf{b}) \rangle + \langle r_i(\theta, \mathbf{b}), H_\theta r_i(\theta, \mathbf{b}) \rangle$$

(H_θ denotes the Hessian) and

$$D_b \mathbf{F}_i(\theta, \mathbf{b}) = \langle D_\theta r_i(\theta, \mathbf{b})^T, D_b r_i(\theta, \mathbf{b}) \rangle + \langle r_i(\theta, \mathbf{b}), D_b \{D_\theta r_i(\theta, \mathbf{b})^T\} \rangle.$$

In general, the above derivatives must be computed during the implementation of the algorithm proposed in Section 2, so no extra computational complexity is incurred in estimating Γ and Λ . The matrices Γ and Λ can be consistently estimated by their sample

counterparts,

$$\begin{aligned}\hat{\mathbf{\Gamma}}_n &= n^{-1} \sum_{i=1}^n \langle \dot{\mathbf{f}}_i(\hat{\mathbf{b}}_n), \dot{\mathbf{f}}_i^T(\hat{\mathbf{b}}_n) \rangle, \\ \hat{\mathbf{\Lambda}}_n &= n^{-1} \sum_{i=1}^n \langle f_i(\hat{\mathbf{b}}_n), \dot{\mathbf{f}}_i(\hat{\mathbf{b}}_n) \rangle \langle f_i(\hat{\mathbf{b}}_n), \dot{\mathbf{f}}_i^T(\hat{\mathbf{b}}_n) \rangle.\end{aligned}$$

The approximate variance of $\hat{\beta}_n(s, t)$ is then $\hat{v}_n(s, t) = \boldsymbol{\phi}(s, t)^T \hat{\mathbf{\Gamma}}_n^{-1} \hat{\mathbf{\Lambda}}_n \hat{\mathbf{\Gamma}}_n^{-1} \boldsymbol{\phi}(s, t)/n$, which can be used e.g. to test whether $\beta^*(s, t) = 0$ for given (s, t) .

5 Simulations

We ran a number of simulations to assess the comparative performance of the proposed method and the common causal functional regression model. The data was generated from the following model:

$$\tilde{y}_i(t) = \int_0^1 \beta_0(s, t) \tilde{x}_i(s) ds + \varepsilon_i(t), \quad (13)$$

$$(x_i, y_i) = (\tilde{x}_i \circ w_{0i}^{-1}, \tilde{y}_i \circ w_{0i}^{-1}), \quad (14)$$

with slope $\beta_0(s, t) = 5e^{-50\{(s-.4)^2 + (t-.6)^2\}} I\{s \leq t\}$, covariates $\tilde{x}_i(s) = z_i e^{-30(s-.4)^2}$ with z_i i.i.d. $N(1, .1^2)$, and error term $\varepsilon_i(t) = u_i \sin(4\pi t)$ with u_i i.i.d. $N(0, .05^2)$. As warping functions we used the one-dimensional exponential family $w_{0i}(t) = (e^{a_i t} - 1)/(e^{a_i} - 1)$, with a_i s equally spaced between -1 and 1 . This is referred to as “Model 1” in Table 1. Ten random pairs (x_i, y_i) are shown in Figure 2 for illustration. The effect of the regression model is, roughly, to shift the peaks from around .4 to around .6.

The performance of an estimator can be evaluated from two different perspectives: the goodness of the estimation of β and the goodness of the prediction of new y s. As measure of estimation error we use the integrated squared error $\text{ISE}(\hat{\beta}) = \iint \{\hat{\beta}(s, t) - \beta(s, t)\}^2 ds dt$, and as measure of prediction error we use the mean squared prediction error $\text{MSPE}(\hat{\beta}) = N^{-1} \sum_{i=1}^N \|y_{n+i} - \hat{y}_{n+i}\|^2$, where $y_{n+1}, \dots, y_{n+N}$ are N additional observations generated from model (13)-(14) but without error term (we used $N = 200$). It is also of interest to study the predicting ability of the estimators for observations that follow model (13)-(14) but whose warping functions fall outside the range of the training data. To this end, we generated samples with warping parameters a_i equally spaced between -1 and 1 for estimation but between 1.5 and 2.5 for prediction. This is referred to as “Model 2” in Table 1.

Figure 2: Simulated data. Ten sample curves simulated from Model 1. (a) Covariates; (b) responses.

(n,k)	Estimation error (Model 1)				Prediction error (Model 1)				Prediction error (Model 2)			
	Optimal		AICC		Optimal		AICC		Optimal		AICC	
	Dynamic	Classical	Dynamic	Classical	Dynamic	Classical	Dynamic	Classical	Dynamic	Classical	Dynamic	Classical
(50,5)	.232	.737	.275	2.08	.776	.307	.863	1.58	5.58	35.4	6.49	66.5
	(.002)	(.0005)	(.005)	(.009)	(.009)	(.003)	(.009)	(.018)	(.044)	(.412)	(.016)	(.306)
(100,5)	.223	.737	3.20	3.77	.774	.257	2.25	.469	5.76	35.7	7.57	45.4
	(.002)	(.0003)	(.040)	(.013)	(.007)	(.001)	(.030)	(.003)	(.035)	(.335)	(.055)	(.179)
(100,10)	.223	.735	.409	2.67	.803	.091	.993	.688	5.18	42.8	6.67	62.4
	(.002)	(.0003)	(.025)	(.005)	(.007)	(.001)	(.030)	(.003)	(.044)	(.300)	(.042)	(.140)
(200,10)	.211	.736	3.86	3.99	.796	.066	1.65	.229	5.15	41.4	5.94	51.4
	(.001)	(.0003)	(.063)	(.015)	(.005)	(.0006)	(.021)	(.002)	(.035)	(.209)	(.050)	(.150)

Table 1: Simulation Results. Mean estimation errors and mean prediction errors of dynamic and common causal regression estimators (Monte Carlo standard errors in parenthesis).

As basis functions for the regression slope we used a truncated-product cubic B-spline basis with k equally spaced knots in $(0, 1)$. The values $k = 5$ and $k = 10$ were considered, for which the number of basis functions were $p = 66$ and $p = 141$, respectively. As warping family for the dynamic estimator we used cubic Hermite splines with a single knot at $\tau_1 = .5$. The smoothing parameter λ was chosen by AICC for both estimators. Specifically, we minimized $\text{AICC}(\lambda)$ over λ s of the form $10^{-\nu}$ with ν in the grid $\{1, 1.25, 1.50, \dots, 5\}$. The minimum $\text{AICC}(\lambda)$ was attained strictly inside the range $(10^{-5}, 10^{-1})$ in all cases, so there was no spurious bias due to truncation of the range of λ .

The mean $\text{ISE}(\hat{\beta})$ and the mean $\text{MSPE}(\hat{\beta}) \times 10^3$, based on 500 replications of each scenario, are reported in Table 1. We also included in Table 1 the smallest estimation and prediction errors attainable for the optimal λ . This allows us to evaluate the ability of the

AICC criterion to select a $\hat{\lambda}$ close to the optimum. As expected, Table 1 shows that the classical estimator cannot provide accurate prediction of new y s and accurate estimation of β simultaneously. We found that for all the simulated samples, the classical estimator invariably attains the lowest prediction error at the smallest λ in the grid (for which $\hat{\beta}$ is so irregular that it does not show any resemblance to the true β), while the smallest estimation error is attained at the largest λ in the grid (for which $\hat{\beta}$ is practically zero). In other words, the low MSPE values that the classical estimator attains for “Model 1” are entirely due to undersmoothing, and are attained by essentially meaningless $\hat{\beta}$ s. In contrast, the dynamic regression estimator attains low prediction error and low estimation error simultaneously. The predicting ability of the dynamic estimator is most clear for “Model 2”, for which the MSPE values are an order of magnitude smaller than those of the classical estimator.

6 Application: Lip Movement Data

In this section we apply the dynamic causal regression method to the data of Malfait and Ramsay (2003) shown in Figure 1. As explained in Section 1, the goal is to predict lip acceleration using the EMG curves of lip neural activity. This data is hard to analyze for a number of reasons: the curves have sharp and localized features; the first EMG spike occurs near 0; there is a lot of time variability; and there are only 29 curves in the sample after removing 3 obvious outliers. Our strategy is then to model β using a relatively large number of well-localized basis functions: we used truncated-product linear B-splines with 10 equispaced knots on each axis, which makes a total of $p = 64$ basis functions. As warping functions we used cubic Hermite splines with knots $\tau = (0, .05, .2, .4, .5, .69)$. We also tried other knot sequences but the results did not change much; what is most important for this data is that *some* kind of warping be used, regardless of the exact family of warping functions.

In this case the AICC criterion is not very useful for choosing the penalization parameter λ , because among λ s with $\text{df}(\lambda) < n$, $\text{AICC}(\lambda)$ always decreases as λ increases. The same behavior was observed for other choices of spline order and number of knots. The likely reason for this behavior, besides the small sample size, is that *after synchronization* there is little variability left to explain. Most of the predictive ability of the explanatory curves resides in their dynamics, not in their amplitude. So we chose $\lambda = 10^{-3}$ subjectively. The resulting warped curves are shown in Figure 3 and the estimated regression slope in Figure 4.

We see that dynamic regression does a good job at aligning these curves. The features of both explanatory and response curves emerge more clearly in Figure 3 than in Figure

Figure 3: Lip Movement Example. (a) Synchronized EMG curves; (b) synchronized lip acceleration curves.

1. In particular the peaks of the EMG curves around times .4 and .5, that were barely discernible in Figure 1(a), are more clear-cut in Figure 3(a). These peaks correspond to the agonistic and antagonistic actions of the lower lip muscle that mark the beginning and the end of the ‘b’.

The estimated regression slope $\hat{\beta}(s, t)$ is shown in Figure 4(a). To facilitate interpretation we plotted in Figure 4(b) the filtered values $\hat{\beta}(s, t)I\{|\hat{\beta}(s, t)| \geq 2\sqrt{\hat{v}(s, t)}\}$, where $\hat{v}(s, t)$ is the variance of $\hat{\beta}(s, t)$ estimated by bootstrap. Contour plots of these surfaces are shown in Figures 4(c) and 4(d). It is not surprising that the sharpest peaks of $\hat{\beta}$ are located on the diagonal near the peaks of the EMG curves, because neural activity has an immediate effect on muscle acceleration. But more interestingly, $\hat{\beta}$ shows some significant off-diagonal values. The most noticeable is the crest at input time s between .2 and .3 and output time $t = .45$, indicating that the neural activity that takes place while the ‘o’ in ‘Bob’ is being produced has an effect on the production of the subsequent ‘b’. This is to be expected, since the position reached by the lower lip for the production of the ‘o’ will determine to a large extent the force necessary to close the lip to produce the second ‘b’. A similar effect was observed by Malfait and Ramsay (2003), but it was less pronounced due to the blurring effect of time variability.

More information about the dynamics of the process can be extracted from the warping functions. The use of interpolating Hermite splines facilitates this because the basis coefficients $\mathbf{c}_i$ satisfy $\mathbf{c}_i = w_i(\boldsymbol{\tau})$, as explained in Section 3, so they can be interpreted as the locations on the curve i of the landmarks corresponding to $\boldsymbol{\tau}$. For our choice of $\boldsymbol{\tau}$, c_{i2} to c_{i5} will roughly correspond to the location of the four characteristic peaks of the EMG curves, as can be seen in Figure 3(a). Thus, the differences $d_{i1} = c_{i3} - c_{i2}$ and

Figure 4: Lip Movement Example. (a) Dynamic regression slope estimator; (b) dynamic regression slope estimator filtered by significance; (c) contour plot of slope estimator; (d) contour plot of filtered slope estimator.

$d_{i3} = c_{i5} - c_{i4}$ indicate the durations of the two ‘b’s and $d_{i2} = c_{i4} - c_{i3}$ the duration of the ‘o’. The pairwise correlations of the differences are $\rho_{12} = -.41$, $\rho_{23} = -.66$ and $\rho_{13} = .26$, with bootstrapped standard errors .19, .10 and .21. Then we see that there is a significant (and negative) correlation between the duration of two consecutive phonemes, but there is no significant correlation between the duration of the two ‘b’s. Of course, more accurate information about the phonemes’ duration would be obtained by identifying the exact peak locations curve by curve; that would be possible for these 29 curves, but would be unfeasible for larger datasets. The advantage of our method is that the parameters $\mathbf{c}_i$ are estimated automatically.

Acknowledgement

This research was partially supported by NSF grant DMS 10-06281.

Appendix

Proof of Theorem 1

The proof employs standard empirical-process techniques [as in e.g. Van de Geer (2000)] and essentially mimics the proof of Theorem 2 in Knight and Fu (2000). Let $\hat{\mathbf{u}}_n = \sqrt{n}(\hat{\mathbf{b}}_n - \mathbf{b}^*)$ and

$$Z_n(\mathbf{u}) = \sum_{i=1}^n \|f_i(\mathbf{b}^* + \mathbf{u}/\sqrt{n})\|^2 + n\lambda_n(\mathbf{b}^* + \mathbf{u}/\sqrt{n})^T \Omega(\mathbf{b}^* + \mathbf{u}/\sqrt{n}).$$

Clearly, it follows from (11) that

$$\hat{\mathbf{u}}_n = \arg \min Z_n(\mathbf{u}) - Z_n(\mathbf{0}),$$

since $Z_n(\mathbf{0})$ is constant in $\mathbf{u}$. A simple Taylor expansion of $f_i(\mathbf{b})$ at $\mathbf{b}^*$ implies that $Z_n(\mathbf{u}) - Z_n(\mathbf{0}) = V_n(\mathbf{u}) + o_P(1)$ with

$$\begin{aligned} V_n(\mathbf{u}) &= \mathbf{u}^T \left\{ \frac{1}{n} \sum_{i=1}^n \dot{\mathbf{f}}_i(\mathbf{b}^*) \dot{\mathbf{f}}_i(\mathbf{b}^*)^T \right\} \mathbf{u} + 2 \left\{ \frac{1}{\sqrt{n}} \sum_{i=1}^n f_i(\mathbf{b}^*) \dot{\mathbf{f}}_i(\mathbf{b}^*)^T \right\} \mathbf{u} \\ &\quad + \lambda_n \mathbf{u}^T \Omega \mathbf{u} + 2\sqrt{n}\lambda_n \mathbf{u}^T \Omega \mathbf{b}^*. \end{aligned}$$

Then, if $\sqrt{n}\lambda_n \rightarrow \ell_0$ as $n \rightarrow \infty$, it follows that $V_n \xrightarrow{D} V$, where V is the stochastic process given by

$$V(\mathbf{u}) = \mathbf{u}^T \mathbf{\Gamma} \mathbf{u} + 2\mathbf{w}^T \mathbf{u} + 2\ell_0 \mathbf{u}^T \mathbf{\Omega} \mathbf{b}^*,$$

with $\mathbf{w} \sim N(\mathbf{0}, \mathbf{\Lambda})$ (note that $E_0\{f_i(\mathbf{b}^*)\dot{\mathbf{f}}_i(\mathbf{b}^*)\} = \mathbf{0}$ because $\mathbf{b}^*$ minimizes $E_0\{\|f_i(\mathbf{b})\|^2\}$.) The Argmin Theorem implies that

$$\hat{\mathbf{u}}_n \xrightarrow{D} \arg \min V(\mathbf{u}) = -\mathbf{\Gamma}^{-1}(\mathbf{w} + \ell_0 \mathbf{\Omega} \mathbf{b}^*)$$

and this completes the proof.


References

- Cai, T. and Hall, P. (2006). Prediction in functional linear regression. *The Annals of Statistics* **34** 2159–2179.
- Crambes, C., Kneip, A., and Sarda, P. (2009). Smoothing splines estimators for functional linear regression. *The Annals of Statistics* **37** 35–72.
- Fritsch, F.N. and Carlson, R.E. (1980). Monotone piecewise cubic interpolation. *SIAM Journal of Numerical Analysis* **17** 238–246.
- Gervini, D. and Gasser, T. (2004). Self-modeling warping functions. *Journal of the Royal Statistical Society (Series B)* **66** 959–971.
- Gervini, D. and Gasser, T. (2005). Nonparametric maximum likelihood estimation of the structural mean of a sample of curves. *Biometrika* **92** 801–820.
- Gohberg, I., Goldberg, S., and Kaashoek, M. A. (2003). *Basic Classes of Linear Operators*. Basel: Birkhäuser Verlag.
- Hall, P. and Horowitz, J. L. (2007). Methodology and convergence rates for functional linear regression. *The Annals of Statistics* **35** 70–91.
- Hastie, T., Tibshirani, R. and Friedman, J. (2009). *The Elements of Statistical Learning. Data Mining, Inference, and Prediction. Second Edition*. Springer, New York.
- Hurvich, C.M., Simonoff, J.S., and Tsai, C.-L. (1998). Smoothing parameter selection in nonparametric regression using an improved Akaike information criterion. *Journal of the Royal Statistical Society (Series B)* **60** 271–293.


- James, G., Wang, J. and Zhu, J. (2009). Functional linear regression that's interpretable. *The Annals of Statistics* **37** 2083–2108.
- James, G.M. (2007). Curve alignment by moments. *The Annals of Applied Statistics* **1** 480–501.
- Jupp, D. L. B. (1978). Approximation to data by splines with free knots. *SIAM J. Numer. Anal.* **15** 328–343.
- Kneip, A., Li, X., MacGibbon, B. and Ramsay, J.O. (2000). Curve registration by local regression. *Canadian Journal of Statistics* **28** 19–30.
- Kneip, A. and Ramsay, J.O. (2008). Combining registration and fitting for functional models. *Journal of the American Statistical Association* **103** 1155–1165.
- Knight, K. and Fu, W. (2000). Asymptotics for lasso-type estimators. *The Annals of Statistics* **28** 1356–1378.
- Liang, H., Wu, H., and Carroll, R. J. (2003). The relationship between virologic and immunologic responses in AIDS clinical research using mixed-effects varying-coefficient models with measurement error. *Biostatistics* **4** 297–312.
- Liu, X. and Müller, H.-G. (2004). Functional convex averaging and synchronization for time-warped random curves. *Journal of the American Statistical Association* **99** 687–699.
- Malfait, N. and Ramsay, J.O. (2003). The historical functional linear model. *Canadian Journal of Statistics* **31** 115–128.
- Müller, H.-G., Chiou, J.-M., and Leng, X. (2008). Inferring gene expression dynamics via functional regression analysis. *BMC Bioinformatics* **9** 60.
- Ramsay, J.O. and Li, X. (1998). Curve registration. *Journal of the Royal Statistical Society (Series B)* **60** 351–363.
- Ramsay, J.O. and Silverman, B. (2005). *Functional Data Analysis (Second Edition)*. Springer, New York.
- Tang, R., and Müller, H.-G. (2008). Pairwise curve synchronization for functional data. *Biometrika* **95** 875–889.
- Tang, R., and Müller, H.-G. (2009). Time-synchronized clustering of gene expression trajectories. *Biostatistics* **10** 32–45.

- Van de Geer, S. (2000). *Empirical Processes in M-Estimation*. Cambridge University Press, Cambridge, UK.
- Yao, F., Müller, H.-G. and Wang, J.-L. (2005). Functional linear regression analysis for longitudinal data. *The Annals of Statistics* **33** 2873–2903.
- Wahba, G. (1990). *Spline Models for Observational Data*. CBMS-NSF regional conference series in applied mathematics, SIAM, Philadelphia.
- Wang, K. and Gasser, T. (1999). Synchronizing sample curves nonparametrically. *The Annals of Statistics* **27** 439–460.
- Wu, H. and Liang, H. (2004). Backfitting random varying-coefficient models with time-dependent smoothing covariates. *Scandinavian Journal of Statistics* **31** 3–19.
- Wu, S. and Müller, H.-G. (2010). Response-additive regression for longitudinal data. *Unpublished manuscript*. Available at <http://anson.ucdavis.edu/~mueller/rare8.pdf>.


(a)




(b)




(c)



(d)



(a)



(b)

